

AD-A218 619

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY DTIC ELECTE MAR 01 1990			3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution unlimited		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report #6			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Brigham Young University		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Provo, UT 84602		7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217-5000			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N000014-88-K-0115	
8c. ADDRESS (City, State, and ZIP Code) 800 No. Quincy Street Arlington, VA 22217-5000		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	
		TASK NO.		WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification) Enantiomeric Recognition of Organic Ammonium Salts by Chiral Dialkyl-, Dialkenyl- and Tetramethyl-Substituted Pyridino-18-Crown-6 Ligands					
12. PERSONAL AUTHOR(S) J. S. Bradshaw, C. Y. Zhu and R. M. Izatt					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) Feb. 20, 1990	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Complexation of the enantiomeric forms of various organic ammonium salts by chiral pyridino-18-crown-6 ligands 1-8 (Figure 1) has been studied by the temperature dependent ¹ H NMR technique. ^{1,2} At low temperatures, the peaks in the ¹ H NMR spectra of the complexes attributable to the hydrogen atoms on the CH ₂ groups attached to the pyridine rings (centered at about δ 4.5), or the methyl hydrogens in the cases of 7 and 8, separated into two peaks of equal intensities. The low temperature peak separations were 40-140 Hz. At high temperatures, the appearance of a single peak is caused by a fast intermolecular or intramolecular face to face guest exchange. The kinetic parameters for the dissociation of these complexes were calculated as reported. ^{1,2} Table I shows the coalescence temperatures (T _c) and ΔG_c^\ddagger for the (continued on back)					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Harold Guard			22b. TELEPHONE (Include Area Code) (202) 696-4409		22c. OFFICE SYMBOL

dissociation of the complexes of 1-8 with various chiral organic ammonium salts. The majority of the data are for complexes of these chiral ligands with the hydrogen perchlorate salts of (R)- and (S)- α -(1-naphthyl)ethylamine (A).

It is evident from these differences in the ΔG_c^\ddagger values in Table I that these chiral ligands exhibit enantiomeric recognition for chiral-forms of various organic ammonium salts. All the (S,S)-ligands formed kinetically more stable complexes with the (R) than with the (S)- form of A. As expected, complexes of the (R,R)- and (R,R,R,R)-ligands with the (S)- form of A were more stable kinetically than those with the (R)- form. The degree of recognition was similar in all ligand-A complexes as shown by the $\Delta\Delta G_c^\ddagger$ values being 0.7 to 0.9 kcal/mol except for 1, 2 and 4 where $\Delta\Delta G_c^\ddagger$ values were 1.1, 1.3 and 1.6 kcal/mole, respectively. An X-ray crystal study of the complexes of (S,S)-1- with both (R) and (S)-A showed that the methyl groups on the chiral carbons of (S,S)-1 interact sterically with one of the naphthylene hydrogens of A in the (S,S)-1-(R)-A complex.¹ We had hoped that larger alkyl groups attached to the chiral centers, such as the sec-butyl groups of 5, would cause even greater enantiomeric recognition. This was not the case. It is possible that the larger substituents cause steric interactions in both sets of complexes.

The reasons for the observed enantiomeric recognition by these ligands for the other chiral salts are not so clear. Ligand 7 showed excellent recognition of the (S)-form of the hydrogen perchlorate salt of α -phenylethylamine (B) over the (R)-form by 2.2 kcal/mol. Even though high recognition values were not evident for the complexes of 5 with the enantiomers of A, ligand 7 did show good recognition for the (R)-form of the hydrogen perchlorate salt of 2-amino-2-phenylethanol (D) over the (S)-form by 1.6 kcal/mol.

1. Davidson, R. B.; Bradshaw, J. S.; Jones, B. A.; Dalley, N. K.; Christensen, J. J.; Izatt, R. M.; Morin, F. G.; Grant, D. M. *J. Org. Chem.* 1984, **49**, 353.
2. Bradshaw, J. S.; Thompson, P. K.; Izatt, R. M.; Morin, F. G.; Grant, D. M. *J. Heterocyclic Chem.* 1984, **21**, 897.

Figure 1. Chiral Pyridino-18-Crown-6 Ligands




Table I. Free Energies of Activation, ΔG_c^\ddagger values (kcal/mole) in CD_2Cl_2 ^a for the Interaction of Chiral Macrocyclic Ligands with Chiral Alkyl Ammonium Salts

Ligand	value ^a	(R)-A ^b	(S)-A ^b	(R)-B ^b	(S)-B ^b	(R)-C ^b	(S)-C ^b	(R)-D ^b	(S)-D ^b
(S,S)-1 ^c	T _c , °C	12	-19			-25	-36		
	ΔG_c^\ddagger	13.4	12.3			12.1	11.8		
(S,S)-2 ^d	T _c , °C	11	-35			-21	-45		
	ΔG_c^\ddagger	13.3	12.0			11.9	10.8		
(S,S)-3 ^c	T _c , °C					-33	-28		
	ΔG_c^\ddagger					11.5	11.6		
(S,S)-4 ^c	T _c , °C	-56	-86			-40	-73		
	ΔG_c^\ddagger	10.3	8.7			11.3	10.0		
(R,R)-5	T _c , °C	-20	5	-50	-15	-42	-48	-30	-39
	ΔG_c^\ddagger	12.5	10.3	10.8	12.1	11.3	10.9	12.6	11.0
(S,S)-6	T _c , °C	29	10						
	ΔG_c^\ddagger	14.2	13.3						
(R,R,R,R)-7	T _c , °C	3	27	-21	25				
	ΔG_c^\ddagger	13.4	14.3	12.1	14.3				
(R,R,R,R)-8	T _c , °C	-52	-43	-38	-46				
	ΔG_c^\ddagger	10.5	11.2	11.2	10.6				

^aA Varian Gemini-200 spectrometer was used to record all ¹H NMR spectra. Equimolar amounts of ligand and salt were dissolved in CD_2Cl_2 . The hydrogens on the CH₂ next to the pyridine ring were used as the ¹H NMR probe for all complexes of 5 and 6 and the methyl hydrogen atoms for 7 and 8. T_c = coalescence temperature. ΔG_c^\ddagger values were ± 0.2 . ^bA = the hydrogen perchlorate salt of (R)- or (S)- α -(1-naphthyl)ethylamine; B = hydrogen perchlorate salt of (R)- or (S)- α -phenylethylamine; C = the hydrogen perchlorate salt of methyl phenylalaninate; D = the hydrogen perchlorate salt of (R) or (S)-2-amino-2-phenylethanol. ^cData for 1, 3, 4 are taken from reference 1. ^dData for 2 were 13 and 12.3 kcal/mole in Reference 2. The measurements were repeated on the Gemini 200 to give the values shown.

By
Distribution/
Availability Codes
Avail and/or
Special

A-1